

The Significance of Palynology in Socio-economic Development in Nigeria.

Ajipe, J.O.

Department of Biological Science
School of Science
Yaba College of Technology
Yaba, Lagos, Nigeria.

Adebayo, M.B.

Department of Botany
Faculty of Science
University of Lagos
Akoka, Lagos, Nigeria.

Corresponding E-mail: johnolusgy@yahoo.com

ABSTRACT

Palynology has become a well-established research tool leading to enormous breakthroughs in the world of science. As a result, Palynology has integrated perfectly into multidisciplinary programmes, such as Melissopalynology, Aeropalynology, Forensic palynology, Agricultural palynology, Fossil palynology, Morphographic palynology (pollen morphology, pollen productivity), Latropalynology, Palaeopalynology including many other areas. Through these multidisciplinary programmes, Palynology has found various applications in Archaeology, Geology, Criminology (Forensic science), Allergology in medicine, Plant taxonomy, Stratigraphy, Paleoecology, Paleobotany and Petroleum industry. The term palynomorph includes both pollen of spermatophytes, spores of fungi, ferns, and bryophytes, as well as other organic-walled microfossils such as dinoflagellates and acritarchs. Pollen grains can be studied for comparative morphological data, clues to unexpected aspects relating to breeding systems, pollination biology and hybridization. This can bring about a better understanding of the whole biology of the group under investigation. Forensic palynology deals with the use of pollen and spores in disproving or proving relationships between people, crime scenes and objects so as to resolve civil or criminal cases. In recent petroleum researches, palynology has become an important tool in resolving many age and facies correlation problems. Palynologists are evolutionarily oriented botanists working ultra-structurally, especially those routinely using Transmission Electron Microscopy (TEM). They are perfectly trained, highly experienced scientists who regularly bring solutions to socio-economic problems. It is in view of this that we present an overview of the various contributions and the significance of palynology, and how it could be further harnessed in bringing about socio-economic development in Nigeria.

Keywords: Melissopalynology, Aeropalynology, Archaeology, Allergology, Paleobotany.

Aims Research Journal Reference Format:

Ajipe, J. O. & Adebayo, M. B. (2018): The Significance of Palynology in Socio-economic Development in Nigeria. *Advances in Multidisciplinary & Scientific Research Journal*. Vol. 4. No.1, Pp51-58

1. INTRODUCTION

Palynology is the study of microscopic bodies generally known as palynomorphs; pollen and spores and certain other microscopic sized structures, either of plant or of uncertain origin. These other structures include acritarchs, dinoflagellates and their cysts (dinocysts), algal spores and fungal spores. The diameters of palynomorphs fall within the range of 5µm-500µm. Thus, "Palynology" can also be defined as the study of organic-walled microfossils (Erdtman, 1969). The term "Palynology" was coined by Hyde and William (1944) and since then, it has become a new sub-division of Botanical science with different applications. At the beginning, Palynology was confined to the study of the morphology of pollen and spores alone. Many people think of pollen only in terms of hay-fever allergies and human health, but its story is really much more interesting. Pollen is the male gametophyte or the microgametophytes of seed plants that produces the male gametes. Some of these pollen grains with features such as bilateral air bladders, microscopic sizes are produced by many angiosperms (for example, conifers) which help during airborne transport of the grains. The angiosperms (flowering plants), produce some pollen with characteristic hook-like structures on the exine wall that allows it to become attached to the legs of insect pollinators.

Spore is usually a haploid and unicellular structure, which is a unit of sexual or asexual reproduction, produced by seedless plants during meiosis in the sporangium of a diploid sporophyte (Erdtman, 1971). Hence, spores and pollen grains are reproductive propagules and play a paramount role in the life history of land plants. The exine walls of pollen and spores are made up of a complex chemically inert biological polymer known as Sporopollenin. It is a very stable chemical compound which makes the pollen and spores to be well preserved in soil and sediments. Present on the wall of the exines are apertures surrounded with aperture membranes. There have been some dispute as to whether or not such membranes are always exinous, for over a century (Wodehouse, 1935). In seedless plants; ferns, the sporophyte generation produces single-celled spores that give rise to the haploid gametophyte generation. A single type of spore is produced in homosporous plants, whereas two spore types are produced in heterosporous plants. The microspores develop by meiotic division to form microgametophytes, and the megaspores also undergo meiotic division to form the megagametophytes. The gametophytes of most seedless plants are multicellular and proliferate outside the spore wall during development. By comparison, all seed plants (gymnosperms and angiosperms) are heterosporous, and pollen represents the microgametophyte. Pollen grains consist of only three to several cells, and these remain contained within the microspore wall, within which they have developed (Osborn, 1997).

The study of pollen grain in the light of modern science started in the 19th century, and as of today, the study of pollen and spores occupy a focal point due to the fact that unique "morphoform" of their structures have been the fundamental factor that served to shape and widen the scope of Palynology. Erdtman and Gunnar (1921), described pollen as a tool for study of the Quaternary vegetation and climate change. The name 'palynogram' was introduced by Erdtman (1951, 1952) to describe a standardized illustration of the morphology of a pollen grain or spore. Additional insight may be obtained into evolution, from the aspect relating to morphological data of fossil pollen and the fossil record. Analysis of fossil pollen grains is the most important approach to reconstruction of past flora, vegetation and environment (Faegri and Iversen, 1989). Palynology is divided into two broad fields, Paleopalynology and Neopalynology. Paleopalynology deals with the study of palynomorphs (including micro fossils) found in the past environment, while Neopalynology is concerned with extant palynomorphs (including the extant pollen). Palynology is becoming increasingly important in basin analysis worldwide. Within the past quarter century palynology has become an increasingly important component of archaeological research. Applications have included elucidation of site and room functions, ceremonial and medicinal practices, prehistoric diet and food preparation, correlative construction and chronologies, human modification of the local environment and the nature, magnitude and duration of climatic perturbations, particularly as related to human demography and subsistence strategies (Hevly, 1981).

1.1 Socio-Economic Development

Socio-economic development is best regarded as the relationships or interactions between economic activity and social life. The process of social and economic development in a society depends largely on the continuous improvement in the wellbeing and in the standard of living of the people. Everyone agrees that development is necessary, and everyone wants it, for instance; Academicians, politicians and economics all agree that development improves the standard of living people. Jan Drewnowski (1966), defined development as a process of qualitative change and quantitative growth of the social and economic reality which we can call either society or economy. Socio-economic development in Socio-economic thematic area aims to ensure that Africans meet their basic needs that are essential to live a life of dignity, which come in forms of access to quality education, better and acceptable healthcare, decent and good housing, safe drinking water and good sanitation, as well as equitable distribution of a nation's wealth. Socio-economic development is measured with indicators such as Gross domestic products (GDP), life expectancy, literacy and levels of employment. Mabogunje (1980) stated that there are four main ways in which the term development has been used: development as progress in the rate of economic growth, as in modernization, as rapid growth in distributional justice and as socio-economic transformation. Changes in less-tangible factors such as personal safety and freedom from fear of physical harm, and the extent of participation in civil society will also boost socio-economic development in our societies. Hence, new technologies, changes in laws, changes in the physical environment and ecological changes are some of the causes of socio-economic impacts.

1.2 Current state of the Nigerian economy

Vanguard newspaper on 8 May, 2017, described the state of the Nigerian economy at half way marked. From the recent data, we can infer that the Nigerian economy is weak and sick. Of course, the cause of this has been attributed to corruption, mismanagement and misappropriation of funds, but truth still remains that Nigeria has been a mono-economic country; more than ninety percent of the nations' gross income is from the oil sector. The Nigerian economy as at present can still be regarded hitherto as a mono-economy. The oil sector (oil and gas) provided up to 95% of Nigeria's foreign exchange earnings and 80% of its budgetary revenues. Due to fall in price of oil in the world market, Nigeria is now experiencing **Economic recession**. According to Sowunmi, in a public lecture (2016), **Economic recession**: "... slowdown in **economic** activity over the course of a normal business cycle." (Wikipedia);

Economic depression has also been defined as a sustained, long-term downturn in economic activity, which has led to more severe downturn than an *economic recession...* (Wikipedia). Now, can we continue to look on while the state of the economy of our “**Dear Nation....NIGERIA**” crashes and pretend that all is well?

1.3 Solutions to this problem

The issue of economic recession has always been a major discussion even during the regime of President Goodluck Jonathan, however, all efforts by the government so far are yet to produce major results. Although President Muhammadu Buhari promised to change the state of the Nigerian economy when he was voted into power, its three years now and promises of change still lie in the air, thus becoming a mirage. Nonetheless, as long as our leaders and everyone are committed to the socio-economic development of Nigeria, we can still hope for a better tomorrow. We hold that there is a need for a shift in the Nigerian economy from a mono-economy to a multi-economic system. Thus, the need for diversification of the economy. The call to diversify the economy was sounded as far back as 1975 but it was unheeded until it became imperative now. The advantage of diversifying include decrease in importation, boost of exportation; being self-reliant in what is consumed; increase in the number of industries among others. The effect of these is that there would be reduction in level of unemployment; socio-economic development; and enhanced quality of life for average citizen (Sowunmi, 2016). Now, why is **Palynology** significant in socio-economic development?

2. HISTORY OF PALYNOLOGY IN NIGERIA

Palynology began in Nigeria in the 1960s when Sowunmi started pioneering Paleo-ethnobotanical and environmental archaeological studies in Nigeria. She set up the first Palynology Laboratory in 1968 at the Archaeology Unit of African Studies (now Department of Archaeology and Arthropology). Today the laboratory has over 3000 references pollen slides of present-day Nigerian and other West African plants (Oyelaran et al., 2016). Since then, a lot of work has been done in Palynology, particularly the study and the reconstruction of the past vegetation of our environment in Nigeria.

Sowunmi (1973) conducted the first Palynological studies when she described the pollen grains of 150 Nigerian woody plants, comprising over 60% of the recognised climatic indicator species of the forest such as primary and secondary rain forests, and swamp forests, and savanna, such as derived forest in the Guinea and Sahel zones, these being the major vegetation zones in Nigeria. Her work was published in *Grana* 13: 145-186, 1973. The paper is the first in a series on the pollen flora of Nigeria. The plants includes; *Lannea microcarpa*, *Mangifera indica* which are of the family Anacardiaceae, *Dracaena arborea* of the family Agaceae, *Annona senegalensis* of the family Annonaceae and others. She compared her findings with those of earlier investigators and she concluded that Intra-specific variations were observed among the morphological structures of the pollen grains and could be used to provide additional characters for use in plant taxonomy. She stated that the pollen could serve as a reference material for pollen analysis of Quaternary deposits.

Sowunmi (1976) continued her laudable contributions to Palynology when she pioneered the analysis of honey pollen from the forest and savanna zones of Nigeria (Melissopalynological investigation). She conducted the pollen analyses of eight honey samples collected from the Guinea savanna, derived savanna, and the dry lowland rain forest. From the study, Sowunmi was able to ascertain the botanical and the geographical origins of some Nigerian honey samples, alongside with the ecology of the area concerned. Agwu and Akanbi (1985), based on the outcome of their palynological research on honey samples warned that the unscientific and traditional agricultural practices are threatening the flora of several parts of Nigeria with increased uncontrolled and indiscriminate destruction. They gave the warning because they were able to ascertain that the vegetation serves as the source of honey. Njokuocha (2006) reported the presence of airborne pollen grains in Nsukka, Nigeria after sampling at two different heights (1.8 m and 15 m) from February 1993 to January 1994. He stated that twenty-six plant families (40 genera) were identified at the lower sampling height, while thirty-eight families (58 genera) were identified at the height of 15 m. The result of his work revealed that: (1) There are differences in pollen concentration at various source heights; (2) There are three periods of varying pollen abundance, and (3) The period of highest pollen concentration is the late rainy – early dry season/Harmattan (September to December), while the low pollen season is during the rainy period (May to August) (Njokuocha, 2006).

Mbah and Amao (2009) investigated the natural foods and feeding habits of the African honey bee *Apis mellifera adansonii* Latrielle, in Zaria, northern Nigeria. They observed and collected 28 plant species visited by the honeybee *Apis mellifera adansonii* Latrielle, using secateurs and plant presses to cut and preserve the specimens, respectively. They also used digital camera to photograph the honeybees on flowers. The plants includes; *Bauhinia grandifolia*, *Acacia ataxacantha*, *Crotalaria falcata*, *Cosmos sulphurens*, *Parkia biglobosa*, *Guiera senegalensis*, *Sida acuta*, *Tridax procumbens*, *Musa sapientum*, and so on. The plants were categorized into 15 families, including; Leguminosae, Asteraceae, Malvaceae, Combretaceae, Mimosaceae, Rutaceae, Musaceae, Portulacaceae, Labiatae, Apocynaceae, Acanthaceae, Caesalpinaceae, Verbanaceae, and Curcubitaceae.

The result of their work based on the categorisation of the plants into 15 families created a pointer to the diverse nature of the natural food sources of these insects (Mbah and Amao, 2009).

Adeonipekun and John (2011) carried out a Palynological investigation of haze dust in Ayetoro-Itele Ota, Southwest Nigeria. The aim of their work was to verify the biological contents of the March 2010 hazy dust in Nigeria as a result of the attendant panic it brought to the public. 3g of dust deposited on a car bonnet was treated chemo-palynologically. The microscopic study of the residue was carried out. It was revealed from their study that the dust was rich in pollen grains, fungal spores and hyphae. There was no record of pteridophyte spore. However, there was record of high proportion of diatoms almost doubling that of the sporomorphs. They concluded that the hazy dust studied contained high proportions of palynomorphs from the Sudan/Guinea and derived savannas as well as lowland rainforest ecozones. Hence, Savanna and derived savanna pollen grains and the Sahara desert freshwater diatoms recovered indicate that the dust was still the harmattan borne by the Northeast Trade winds though coming heavily and lately in March instead of the traditional January (Adeonipekun and John, 2011).


Mbagwu et al. (2009) palynologically studied five species of the family Asteraceae namely *Guternbergia nigriflora*, *Emilia praetensis*, *Vernonia guineensis*, *Lagera pterodonta* and *Chromola odorata*. Results obtained from their work showed that pollen shape is spheroidal in *G. nigriflora*, *E. praetensis* and *C. odorata* while it is elliptic in *V. guineensis* and *L. pterodonta*. The pollen aperture was found to be porate in all except in *L. pterodonta* where it is elliptic. The pollen wall is echinate in all except in *C. odorata* where it is smooth. They also concluded that the differences and similarities in pollen morphology of the investigated species are significant and could be exploited for biosystematics purposes. Hence, applying these variations in pollen morphology to the species investigated showed that species in the family with similar pollen characters are more closely related and thus exhibit interspecies relationships suggesting reasons for them to be in the same family while those with different pollen characters are not very closely related and suggest reasons for them to be as distinct species (Mbagwu et al., 2009).

Adebayo et al. (2012), conducted a research on the Palynology of Bog-1 Well, Southeastern Niger Delta Basin, Nigeria. They assigned a Late Oligocene-Mid Miocene age to the miospores recovered from the sediments of a section of Bog-1 well in the Niger Delta Basin. Zonations were based on the co-occurrence of pantropical stratigraphic markers such as *Zonocostites ramona*, *Retimonocolpites pluribaculatus*, *Retibrevitricolporites protrudens*, *Psilatricolporites crassus*, *Circatricolporites doroensis*, *Retitricolporites irregularis*, *Racemonocolpites hians*, *Pachydermites diederixi*, *Brevicolporites guinetii* and *Proxapertites cursus*. Moreover, the palynological assemblage was dominated by pollen and spores and a total of 119 palynomorphs were identified out of the 152 encountered. Species diversity of above 30 was regarded as rich while a diversity value below 30 was regarded as poor. There was a general occurrence of high species diversity, except at a few levels. They concluded that the diversity of the angiosperm palynoflora, which forms the bulk of the assemblage, suggests a dense lowland vegetation cover during the deposition of the studied sediments in a tidal swamp mangrove (Adebayo et al., 2012).

Chiaghanam et al. (2013) described the Palynology and Palaeoenvironmental Study of Early Campanian to Mid-Maastrichtian Deposits of Udi and Environs in the Anambra Basin, South Eastern Nigeria. They made use of the Palynological assemblages, lithofacies and facies association to determine the age and palaeoenvironment of the Enugu and Mamu Formations of Udi area of the Anambra Basin, South east Nigeria. Palynological analysis were carried out in over twenty samples which yielded spores and pollen grains alongside with marine Dinoflagellates that are of Early Campanian to Maastrichtian for Enugu shale and middle Maastrichtian for Mamu Formation. According to Chiaghanam et al. (2013), the main diagnostic species of spores and pollen recovered includes; *Laevigatosporites ovatus*, *Leiotriletes adriennis*, *Echitriporites trianguliformis*, *Longapertites marginatus* and *Cyathidites minor*. Among the main diagnostic dinoflagellates are *Operculodinium centrocarpum*, *Areoligera senoniensis*, *Spiniferites ramosus*, *Ceratiopsis* spp., *Paleocystodinium* spp., *Dinogymnium accuminatum*, *Spiniferites ramosus* and *Cordosphaeridium inorders*.

Moreover, they concluded that the Palynological analysis and lithofacies association which was used to determine the age and palaeoenvironment of the studied sections are indicative of a deltaic sedimentation in an Anambra basin of a Campanian maastrichtian sedimentary unit that are characterized with transgressive and regressive events as reflected in the abundance and diversity of palynomorphs and dinoflagellates that were recorded. The palynomorphs and dinoflagellates distribution in the unit showed that Enugu shales were deposited in a deeper marine environment than the Mamu Formation and it is an indicative of a drop in sea level, possibly saline water and fluvial processes (fresh-water) in a fresh water swamp/upper deltaic plain (Chiaghanam et al., 2013). Subsequently, Paul et al. (2014) described the comparative study of the pollen morphology of some members of Euphorbiaceae family. From their studies, they concluded that most pollen grains of Euphorbiaceae family are isopolar and radially symmetrical.

Moreover, the variations and similarities in pollen morphological characters within the genera and species are enough to confirm and throw more light on its taxonomic classifications. They made a recommendation that the studies will be useful for identifying the flora and to improve the conservation status of economically important plants as well as act as a reference guide in identifying various species in their respective families (Paul et al., 2014).



(1) *Murraya koenigii*, Polar view (PV)

(2)a. *Talinum triangulare*

(2)b. Pore outlines.



(3) *Elaeis guineensis*



(4) *Ageratum conyzoides*



(5) *Cochorus olitorius*
Equatorial view (EV)

Plates 1 - 5: Photomicrographs of the pollen grains of some Nigerian plants.

3. BRANCHES OF PALYNOLOGY

3.1 Forensic palynology

Forensic palynology is the application and utilization of palynomorph analysis into crime and law. Here, the use of microscopic pollen and spores (also called palynomorphs) is introduced as an activity in science of forensic palynology to solve criminal cases. The various features of pollen and spores such as small size, high variability, and being found on almost any item that has been exposed to or comes in contact with the air make them the ideal forensic trace materials (Milne *et al.*, 2005). There is always transfer of pollen and spores between people, plants, and objects or there may be a combination of contact between these things. Pollen can be isolated from most items typically submitted for forensic examination. These items could be soil, shoes, drugs, clothing and fabrics, ropes and twines, air filters, plant materials such as leaves, stems, as well as human and animal material, such as hair, fur, and stomach contents (Horrocks 2004; Milne *et al.* 2005). Scientists make use of a digital pollen database and Google Earth to link pollen trace evidence to a specific crime scene. The methods used in forensic palynology, involve those used in criminal cases and palynological techniques used by forensic scientists. Due to its prominence in popular media, forensic science is appealing to this current generation of students. Forensic palynology can harness this excitement and focus student attention on subtle aspects of ecosystems such as the role of pollen and spores in plant propagation (Babcock and Warny, 2014). The demonstrations of forensic palynology in crimes such as rape, homicide, genocide, terrorism, drug dealing, robbery, and others constitute its historic published applications (Milne, Bryant, and Mildenhall 2004).

3.2 Aeropalynology

This is the branch of palynology that deals with the study of palynomorphs suspended in the air including those with allergenic effects such as the hay fever. It is the study of pollen grains and spores that are dispersed into the atmosphere. Studies have shown differences in the pollen concentration at various heights and seasons. Wind-pollinated plants are known to typically produce copious amounts of pollen. The abundance of the pollen in the air help in enhancing successful pollinations. Daily observations of the airborne pollen indicate that pollen grains and fern spores are suspended in air turbulence. They are transported over varied distances away from their sources to other areas of the same or different ecological characteristics (Bringfelt et al., 1982). Pollen abundance or concentration has always been lower at wet or rainy seasons and higher at dry season or Harmattan (Njokuocha, 2007). For different plants, flowering time and season vary widely, and the release of airborne pollen is typically inhibited by high humidity or rain. Moreover, the abundance of airborne pollen commonly causes allergic reactions in a large proportion of the human population, for example, Pollinosis, Allergin Rhinitis, or Hay fever, which is elicited when allergen-containing pollen makes contact with the mucous membranes lining the nose, trachea, or bronchi and the cornea of the eye. Allergens leach out of the pollen and bind to immunoglobulin E antibodies. The antibodies are linked to mast cells that release histamine and other inflammatory chemicals, producing allergy symptoms. Ironically, the allergens that induce pollinosis include many of the same compatibility-incompatibility, recognition proteins involved in pollination (Osborn, 1997). In understanding and avoiding hay fever, knowledge of the temporal, seasonal, and environmental aspects of pollen dispersal is also important, and more also, to monitor the risks of pollinosis, the diversity and quantity of various pollen types are assessed by filtering the air throughout the year.

3.3 Application in Medicine and Aeropalynology

As stated earlier, the biological contents of the March 2010 hazy dust in Nigeria deserve verification due to the attendant panic it brought to the public (Adeonipekun and John, 2011). Allergenic pollen produce certain chemicals, most important are histamines. Symptoms of pollinosis include sneezing, conjunctivitis, pneumonia, trachea-bronchitis, sinuses, possibly also migraine headache, uterine pain, dermatitis (small reddish swellings on skin). Possible "culprits" like *Zea mays*, fungal spores e.g. *Cladosporium* sp. have been suspected to trigger allergic reactions. Pollinosis can be treated or avoided by administering anti-histamine drugs and health-promoting pollen tablets because of their nutritive values, esp. proteins and mineral salts, and keeping susceptible persons indoors. Pharmaceutical palynology is the field of palynology that studies and explores the manufacturing of pollen tablets.

3.4 The role of pollen morphology in plant systematics

Pollen morphology is an expression of part of the genome and like any character be it cryptic or macromorphological, it may be useful in some groups for taxonomic studies and less valuable in others (Ferguson, 1985). For about 150 years, comparative pollen morphology has been studied by the likes of Mohl (1835) and Hassell (1842). Erdtman (1952) provided the work which is rightly regarded as the corner stone of modern comparative pollen morphology. His work centred around the study of the acid resistant sporopollenin exine. Characters such as shape, size, apertures, ornamentation and the stratification of the wall have proved so useful in distinguishing pollen grains of different varieties, species, genera, tribes, families and orders. The functional aspects of pollen such as pollination biology and pollen- pistil interaction have also been found to be well understood through pollen morphology. In Nigeria, not much work have been done on pollen morphology.

3.5 Application of Palynology in stratigraphy (stratigraphy palynology)

Stratigraphic palynology is a branch of biostratigraphy that deals with applying geological and evolutionary principals to the understanding of sedimentary sequences and the geological record. The correlation among sedimentary sequences of any age and the provision of the chronological control for these sedimentary sequences involve the utilization of palynomorphs alongside with their identification, distribution, and abundance. Typically, stratigraphic palynology is applied to pre-Quaternary sediments but also to sediments of all ages. Stratigraphic palynology played a prominent role in petroleum exploration during the mid-twentieth century. The information palynology provides has proven over the years that palynology is an extremely significant parameter for generating new geological models and refining hydrocarbon (oil and gas) exploration researches (Mehrotra *et al.*, 2012). Historically in North-America, the major petroleum companies hosted large palynology laboratories from 1965 to 1980. The ubiquity of palynomorphs permitted the detailed analysis of subsurface stratigraphy necessary for locating petroleum reservoirs. Analysis of fossil pollen grains is the most important approach to the reconstruction of past flora, vegetation and environment (Faegri and Iversen, 1989).

3.6 Application in apiculture (bee-keeping) - Melissopalynology

Melissopalynology is a branch of palynology that deals with the study of pollen grains, spores and other microscopic objects contained in honey samples. The term “melissopalynology” was coined from the specific name of honey bees, “*Apis mellifera*”, and “palynology”. Honey is a viscous, sticky and sweet liquid made from several sweet plant juices, mainly nectar and opened, ripe fruits, as well as honey dew (Crane, 1975). Melissopalynology also began in 1976 in Nigeria at the Palynology Unit of the Department of Archaeology and Anthropology, University of Ibadan, when Sowunmi carried out the first melissopalynological investigation (Sowunmi, 1976). Since then, a lot of work have been done on melissopalynological analysis of Nigerian honey, alongside with the analysis of their physico – chemical components in the various vegetational and geographical zones of Nigeria, by several researchers, and these have greatly helped in determining the botanical and geographical origins of the honey samples through the pollen types that were observed, and ascertaining purity of sample and authenticity of provenance as claimed by marketer. Thus, all the melissopalynological studies that have been carried out in Nigeria have shown that the Nigerian honey samples are characterized by a much diversified pollen flora, alongside with some physical and chemical components. However, there are adulterated forms of honey samples in Nigeria markets, thus, for Nigerian honeys to command respectability both nationally and internationally, efforts should be made to checkmate this. This has begun in University of Ibadan, Nigeria (Ajipe, 2015).

4. CONCLUSIONS

Pollen, being a strong, characteristic and distinguishable unit cell of a flower play important role in Palynology. Palynology is a field with diverse applications in many disciplines including Entomology, Archaeology, Ecology, Biology, Ethnobiology, Medicine, Forensic, Systematics, Taxonomy, Environmental studies, Geoengineering and Human diet. Palynology provides opportunity for collaborations between the academia and the various industries that rely on the knowledge and discoveries from palynological research in driving socio-economic development in Nigeria. Palynologists are evolutionarily oriented Botanists who work ultra-structurally, especially those routinely using Transmission Electron Microscopy (TEM). They are perfectly trained, highly experienced scientists who regularly bring solutions to socio-economic problems. As a result, we recommend that the study of Palynology should be extended to most institutions in Nigeria as this would further amplify the direct and indirect effect of Palynology on the socio-economic state of the nation while also stirring the interest of both undergraduate and postgraduate students in pursuing a career as a Palynologist. Although we do not have many palynologists in Nigeria, Palynology will still continually play its role as a major tool in research and socio-economic development.

REFERENCES

1. Adebayo, O. F., Orijemie, A. E. and Aturamu, A. O. (2012). Palynology of Bog-1 Well, Southeastern Niger Delta Basin, Nigeria. *International Journal of Science and Technology*, 2(4): 214-222.
2. Adekanmbi, O. H. and Ogundipe, O., (2005). Pollen grains of some cultivated plants in Nigeria. *Journal of Science Research and Development*, 10: 101-110.
3. Adeonipekun, P. A. and John. M. (2011). Palynological investigation of haze dust in Ayetoro-Itele Ota, Southwest Nigeria. *Journal of Ecology and the Natural Environment*, 3(14): 455-460.
4. Agwu, C.O.C. and Akanbi, T.O., (1985). A palynological study of honey from four vegetation zones of Nigeria. *Pollen et spores*, 27(3-4): 335- 348.
5. Ajipe, J.O. (2015). A melissopalynological study of honey samples from South Western Nigeria: towards a standardization and quality control of honey in Nigeria. An unpublished M.Sc. dissertation, Botany Department University of Ibadan, Nigeria.
6. Babcock, S. L. and Warny, S. (2014). *Forensic Palynology as Classroom Inquiry*. Routledge Taylor & Francis Group, 51:116–128.
7. Chiaghanam, O.I., Nwozor, K.K., Chiadikobi, K.C., Omoboriwo, A.O., Soronnadi-Ononiwu, C.G., Onuba, L.N. and Ofoma, A.E. (2013). Lithofacies, Palynology and Paleoenvironmental Study of Early Campanian to Mid-Maastrichtian Deposits of Udi and Environs in the Anambra Basin, South Eastern Nigeria. *International Journal of Science and Technology*, 2(6): 453-470.
8. Erdtman, G. (1952). Pollen morphology and plant taxonomy – angiosperms. Stockholm: Almqvist and Wiksell.
9. Erdtman, G. 1960. The acetolysis method - A revised description. *Svensk Botanisk Tidskrift*, 54: 561-564.
10. Erdtman, G. (1969). An introduction to the study of pollen grains and spores. Handbook of palynology, Hafner publishing company. New York. 486pp.
11. Erdtman, G. (1971). Pollen morphology and plant taxonomy. Hafner, New York.
12. Erdtman, O. and Gunnar E. (1921). Pollenanalytische Untersuchungen von Torfmoooren and marinen Sedimenten in Sudwest-Schweden. *Arkiv for Botanik*, 17:10.
13. Traverse, A. (2007). Topics in Geobiology. Paleopalynology, Second edition. 28: ISBN 978-1-4020-5609-3.
14. Ferguson, I.K. (1985). The role of Pollen morphology in plant systematics. *An. Asoc. Pal inol. Leng.*, 2:5-18
15. Faegri, G. & Iversen, J. (1950). *Text-Book of Modern Pollen Analysis*. Copenhagen.
16. Hevly, R. H. (1981). Pollen Production, Transport and Preservation: Potentials and Limitations in Archaeological Palynology. *Journal of Ethnobiology*, 1 (1): 39-54.
17. Jan Drenowski, (1966). Social and Economic Factors in Development, UNRISD, Report Ho. 3, Geneva, Feb., p. 7.
18. Mabogunje, A.L. (1980), op. cit. p. 38.
19. Mbah, C. E., and Amao, A. O. (2009). Natural Foods and Feeding Habits of the African Honey bee (*Apis mellifera adansonii*). *Science World Journal*, 4(1):11-14.
20. Mbagwu, F. N., Chime, E. G. and Unamba, C. I. N. (2009). Palynological studies on five species of Asteraceae. *African Journal of Biotechnology*, 8 (7): 1222-1225
21. Osborn, J. M. (1997). Palynology. In: McGraw-Hill Encyclopedia of Science and Technology, McGraw-Hill, New York. 13: 70-74.
22. Mehrotra, N.C., Shanmukhappa, M., Rupendra, B., Kumar, M., Singh, A., Singh, B. D. and Kapoor, P.N. (2012). Development of Palynology in Fossil Fuel Exploration in India with Emphasis on Recent Significant Contributions from Western-Offshore, Krishna-Godavari Basin and Frontier Areas. *Proceedings of the Indian National Science Academy*, 78(3): 457-473.
23. Njokuocha, R. C. (2006). Airborne pollen grains in Nsukka, Nigeria. *Grana*, 45: 73–80.
24. Oyelara, P.A., Alabi, R.A. and Adeonipekun, P.A. (2016). Human Palaeoecology in Africa. *Essays in honour of M. Adebisi Sowunmi*.
25. Paul, E., Essien, B. C, Idachaba, S. O, Edegb, E. and Tamenku, M. M. (2014). Comparative Study of Pollen Morphology of some members of Euphorbiaceae family. *Standard Research Journal of Agricultural Sciences*, 2(4): 054 – 058.
26. Sowunmi, M.A. (1976). The potential value of honey in Palaeopalynology and Archaeology. *Review of Palaeobotany and Palynology*, 21: 171- 185.
27. Sowunmi, M.A. (1978). Pollen of Nigerian Plants. *Grana*, 13:145-186.
28. Sowunmi, M. A. (2001). Microscopic analysis of Honey. *The Nigerian Field*, 66:125-133.
29. Sowunmi, M.A. (2016). The role of pollen in health care delivery and socio-economic development in Nigeria. (Paper presented at first workshop of the Palynological Association of Nigeria, University of Lagos, 22nd November, 2016.
30. Wodehouse, R.P. (1935). *Pollen Grains*. New York and London.